(1) A sample of helium has a volume of 400 mL at 1.20 atm. What will the volume be at 1.50 atm?

(2) A balloon has a volume of 8.00 L at 47 °C. At what temperature will the balloon have a volume of 7.60 L?

(3) A gas cylinder has a pressure reading of 7.0x10<sup>4</sup> Pa at 350 K. What will the pressure read at 400 K?

(4) A sample of gas has a volume of 500 mL at 3.00 atm and 200 K. What will the pressure be if the sample is expanded to 600 mL at 300 K?

(5) How many moles of carbon dioxide are found in 5.6 L at STP? What is the mass of the carbon dioxide?

(6) How many moles of helium occupy 400 mL at 3.4 atm and 60 °C? What is the mass of helium?

(7) Methane (CH<sub>4</sub>) gas reacts with oxygen to form carbon dioxide gas and water vapour.

(a) Write a balanced chemical equation for this reaction.

- (b) If 448 mL of methane are present at STP, what mass and volume of oxygen are required in the reaction?
- (c) What is the volume and mass of each of the products?

(8) Determine the concentration of a solution containing 0.500 mol hydrochloric acid in 800 mL.

(9) What is the volume if a 0.15 M solution contains 0.24 mol of sodium chloride?

(10) Determine the final concentration if 40 mL of water are added to 60 mL of 0.25 M silver nitrate solution.

| (11) Write a dissociation equation an       | d determine the concentration of each ion in the solution. |
|---|--|
| (a) 0.036 M Na <sub>2</sub> SO <sub>4</sub> | (b) 0.40 M AlCl <sub>3</sub>                               |

| (12) Determine if the follow | wing compounds are solub | ble or insoluble in water. |                |
|------------------------------|--------------------------|----------------------------|----------------|
| (a) AgI                      | (b) SrS                  | (c) $Ca(OH)_2$             | (d) $Na_2SO_3$ |

(13) Write the formula equation, complete ionic equation, and net ionic equation for the reaction between NaI and Pb(NO<sub>3</sub>)<sub>2</sub>.

(14) 150 mL of 0.200 M strontium chloride solution are reacted with 200 mL of silver nitrate solution.

- (a) Write a balanced chemical equation for this reaction.
- (b) What is the concentration of the silver nitrate solution?
- (c) What is the mass of each of the products?

(15) List three properties of acids and three properties of bases.

(16) Complete the following table.

| [H <sup>+</sup> ]      | рН   | рОН  | [OH <sup>-</sup> ]     | acidic or basic? |
|------------------------|------|------|------------------------|------------------|
|                        |      | 3.15 |                        |                  |
| 3.2x10 <sup>-4</sup> M |      |      |                        |                  |
|                        |      |      | 2.8x10 <sup>-6</sup> M |                  |
|                        | 5.05 |      |                        |                  |

(17) Write a balanced equation for each of the following reactions. Classify the reactions.

(a) nitric acid + barium hydroxide  $\rightarrow$ 

(b) sulphuric acid + potassium hydroxide.

(18) Calculate the amount of energy released when 15.0 g of acetic acid freezes.

(19) Calculate the mass of ethanol that can be boiled with  $2.93 \times 10^3$  J of energy.

(20) Calculate the heat energy required to increase the temperature of 100 g of air from 20.0 °C to 80.0 °C.

(21) Calculate the final temperature if 65 J of heat energy is added to 25 g of lead at 45 °C.

(22) How much energy is required to turn 22 g of ice at -12 °C into steam at 114 °C? How many kJ is this?

| (23) Classify each reaction as endothermic or exothermic and determine $\Delta H$ .                        |        |
|--|--------|
| (a) CO (g) + SiO <sub>2</sub> (s) + 590.2 kJ/mol $\rightarrow$ SiO (g) + CO <sub>2</sub> (g); $\Delta$ H = | kJ/mol |

(b)  $2ZnS(s) + 3O_2(g) \rightarrow 2ZnO(s) + 2SO_2(g) + 878.3 \text{ kJ/mol}; \Delta H = ____k \text{J/mol}$ 

- (24) (a) Use the heats of formation to calculate the heat of the following reaction:  $C_3H_8(g) + 5O_2(g) \rightarrow 3CO_2(g) + 4H_2O(g)$
- (b) How much heat would be released by the combustion of 10.000 mol of propane (C<sub>3</sub>H<sub>8</sub>)?
- (25) Write the nuclide symbol for each isotope. State the number of protons, electrons, and neutrons for each isotope. (a) Tellurium–120 (b) Lanthanum–139 (c) Vanadium–50

(26) For each of the following statements, state which type(s) or radiation they describe.

- (a) has the highest penetrating power
- (b) has the same structure as an electron
- (c) has the same structure as a helium nucleus
- (d) can be stopped by a piece of paper

- (e) can be stopped by aluminum foil
- (f) can result in a transmutation
- (g) is energy released from an excited atom
- (h) is a type of particle

(27) Complete each of the following nuclear reactions.

(a) 
$${}^{66}_{29}\text{Cu} \rightarrow {}^{66}_{30}\text{Zn} + \_\_\_\_$$
  
(b)  ${}^{185}_{79}\text{Au} \rightarrow {}^{181}_{77}\text{Ir} + \_\_\_\_$   
(c)  ${}^{209}_{84}\text{Po}^* \rightarrow {}^{209}_{84}\text{Po} + \_\_\_\_\_$   
(d)  ${}^{235}_{92}\text{U} + {}^{1}_{0}\text{n} \rightarrow {}^{94}_{36}\text{Kr} + \_\_\_+ 3{}^{1}_{0}\text{n}$ 

(e) 
$${}^{39}_{17}\text{Cl} \rightarrow {}^{0}_{-1}\text{e} + \_\_\_$$
 (f)  ${}^{100}_{41}\text{Nb} \rightarrow {}^{4}_{2}\text{He} + \_\_\_$ 

(28) The half-life of radium-224 is 3.660 days.

(a) What mass of a 10.0 g sample will remain after 7.32 days?

(b) How long will it take for a 60.00 g sample to decay to 3.750 g?

(29) The half-life of radium-226 is 1599 years.

(a) What mass of a 15.00 g sample will remain after 6396 years?

(b) How long will it take for an 80.00 g sample to decay to 1.250 g?

(30) Sulphur–35 has a half–life of 87.10 days.

(a) What mass of a 64 g sample will remain after 348.4 days?

(b) How long will it take for a 1024 g sample to decay to 4.000 g?

(c) How long will it take for a sample to decay to 12.50% of the original amount?

(d) After 522.6 days, there are 2.00 g of a sample remaining. What was the mass of the original sample?

(31) Name the following compounds.

(a) (b)  $CH_2CH_3$  (b)  $CH_2CH_3$   $CH_3$   $CH_3CH_2CHCH_2CHCH_2CH_3$   $CH_3CH=CHCHCH_3$  $CH_2CH_3$  (c) (d)  $CH_2CH_3$  (d)  $CH_2CH_3$   $CH_2CH_3$   $CH_2CH_2CH_3$   $CH_3C \equiv CCHCHCH_3$   $CH_2CH_2CH_3$   $CH_3C \equiv CCHCHCH_3$   $CH_3C \equiv CH_3CH_3$  $CH_3CH_3CH_3$ 

(32) Draw the following compounds(a) 2,3–Dimethyloctane

(c) 3-methyl-2-heptene

(b) 3,4–Diethylcyclohexene

(d) 5-Ethyl-2-nonyne

(33) Classify each of the compounds and match them with the correct name.



(34) Use SSR to explain what would happen to  $[C_5H_{11}N]$  for each of the following stresses.  $C_5H_5N(g) + 3H_2(g) + 1.4 \text{ kJ/mol} \rightleftharpoons C_5H_{11}N(g)$ (a) increase pressure (b) decrease temperature (c) increase volume

(d) increase [H<sub>2</sub>]

| (35) Write a Kee | q expression for  | each of the                       | following | equilibria |
|------------------|-------------------|-----------------------------------|-----------|------------|
| (a) 2CC          | $O(g) + O_2(g) =$ | $\Rightarrow$ 2CO <sub>2</sub> (g | )         |            |

(b) 4HCl (g) +  $O_2$  (g)  $\rightleftharpoons$  2H<sub>2</sub>O (l) + 2Cl<sub>2</sub> (g)

(36) Write a Keq expression for each of the following equilbria. Determine the value of Keq. Does the equilibrium favour the products or the reactants?

(a)  $2H_2O(l) + 2F_2(g) \rightleftharpoons 4HF(g) + O_2(g)$ At equilibrium,  $[F_2] = 0.160 \text{ M}, [HF] = 1.20 \text{ M}, \text{ and } [O_2] = 0.200 \text{ M}$  (b)  $2CH_4$  (g)  $\rightleftharpoons C_2H_2$  (g) +  $3H_2$  (g) At equilibrium, a 10.0 L container holds 2.00 mol CH<sub>4</sub>, 0.400 mol C<sub>2</sub>H<sub>2</sub>, and 6.00 mol H<sub>2</sub>

## Answers:

(1) 320 mL (2) 304 K (3)  $8.0x10^4$  Pa (4) 3.75 atm (5) 0.25 mol and 11 g (6) 0.050 mol and 0.20 g

(7) (a)  $CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O_{(b)}$  (b) 0.896 L and 1.28 g  $O_2$  (c) 0.448 L and 0.882 g  $CO_2$ , 0.896 L and 0.721 g  $H_2O_2$ 

(8) 0.625 M (9) 1.6 L (110) 0.15 M

(11) (a)  $Na_2SO_4 \rightarrow 2Na^+ + SO_4^{2-}$ ,  $[Na^+] = 0.072$  M,  $[SO_4^{2-}] = 0.036$  M (b)  $AlCl_3 \rightarrow Al^{3+} + 3Cl^-$ ,  $[Al^{3+}] = 0.40$  M,  $[Cl^-] = 1.2M$ 

(12) (a) insoluble (b) soluble (c) insoluble (d) soluble

(13) formula equation: 2NaI (aq) + Pb(NO<sub>3</sub>)<sub>2</sub> (aq)  $\rightarrow$  2NaNO<sub>3</sub> (aq) + PbI<sub>2</sub> (s) complete ionic equation: 2Na<sup>+</sup> (aq) + 2I<sup>-</sup> (aq) + Pb<sup>2+</sup> (aq) + 2NO<sub>3</sub><sup>-</sup> (aq)  $\rightarrow$  2Na<sup>+</sup> (aq) + 2NO<sub>3</sub><sup>-</sup> (aq) + PbI<sub>2</sub> (s) net ionic equation: Pb<sup>2+</sup> (aq) + 2I<sup>-</sup> (aq)  $\rightarrow$  PbI<sub>2</sub> (s)

(14) (a) (a) 
$$SrCl_2 + 2AgNO_3 \rightarrow Sr(NO_3)_2 + 2AgCl$$
 (b)  $[AgNO_3] = 0.300 \text{ M}$  (c) 6.35 g  $Sr(NO_3)_2$  and 8.60 g AgCl

(15)

| Acid                                      | Base                                   |
|---|--|
| dissociate to give H <sup>+</sup> ions    | dissociate to give OH <sup>-</sup> ion |
| pH<7.0                                    | pH>7.0                                 |
| taste sour                                | taste bitter                           |
| react with metals to produce hydrogen gas | feel slippery                          |
| pH paper turns red/orange                 | pH paper turns blue/green              |
| phenolphthalein $\rightarrow$ colourless  | phenolphthalein $\rightarrow$ pink     |
| bromothymol blue $\rightarrow$ yellow     | bromothymol blue $\rightarrow$ blue    |
| cabbage juice $\rightarrow$ pink          | cabbage juice $\rightarrow$ blue       |

(16)

| [H <sup>+</sup> ]       | рН    | рОН   | [OH <sup>-</sup> ]      | acidic, basic,<br>or neutral? |
|-------------------------|-------|-------|-------------------------|-------------------------------|
| 1.4x10 <sup>-11</sup> M | 10.85 | 3.15  | 7.1x10 <sup>-4</sup> M  | basic                         |
| 3.2x10 <sup>-4</sup> M  | 3.49  | 10.51 | 3.1x10 <sup>-11</sup> M | acidic                        |
| 3.5x10 <sup>-9</sup> M  | 8.45  | 5.55  | 2.8x10 <sup>-6</sup> M  | basic                         |
| 8.9x10 <sup>-6</sup> M  | 5.05  | 8.95  | 1.1x10 <sup>-9</sup> M  | acidic                        |

(17) neutralization (a)  $2HNO_3 + Ba(OH)_2 \rightarrow Ba(NO_3)_2 + 2H_2O$  (b)  $H_2SO_4 + 2KOH \rightarrow K_2SO_4 + 2H_2O$ 

(20) 5.97x10<sup>3</sup> J

(18) 2.88x10<sup>3</sup> J

 $x10^3 J$  (19) 5.00 g

(21) 65 °C (22)  $6.7 \times 10^4 \text{ J}, 67 \text{ kJ}$ 

(23) (a) endo;  $\Delta H$ = +590.2 kJ/mol (b) exo,  $\Delta H$ =-878.3 kJ/mol

(24) (a) -2043.8 kJ/mol (b) -20438 kJ

(25) (a)  $\frac{120}{52}$  Te 52 p, 52 e, 68 n (b)  $\frac{139}{57}$  La 57 p, 57 e, 82 n (c)  $\frac{50}{23}$  V 23 p, 23 e, 27 n

(26) (a) gamma (b) beta (c) alpha (d) alpha(e) beta (f) alpha/beta (g) gamma (h) alpha/beta

(27) (a)  ${}^{0}e$  (b)  ${}^{4}He$  (c)  ${}^{0}\gamma$  (d)  ${}^{139}Ba$  (e)  ${}^{39}Ar$  (f)  ${}^{96}Y$ -1 2 0 56 18 39

(28) (a) 2.50 g (b) 14.64 days (29) (a) 0.9375 g (b) 9594 years

(30) (a) 4.0 g (b) 696.8 days (c) 261.3 days (d) 128 g



(33) (a) #7, alcohol (b) #3, amine (c) #1, carboxylic acid (d) #5, ketone (e) #2, aldehyde and alkyl halide (f) #8, amide (g) #4, ether (h) #6, ester

(34)(a) S: increase pressure S: right: R: increase  $[C_5H_{11}N]$ (c) S: increase volume S: left: R: decrease  $[C_5H_{11}N]$  (b) S: decrease temperature S: left R: decrease  $[C_5H_{11}N]$ (d) S: increase  $[H_2]$  S: right R: increase  $[C_5H_{11}N]$ 

(35) (a) 
$$Keq = \frac{[CO_2]^2}{[CO]^2[O_2]}$$
 (b)  $Keq = \frac{[Cl_2]^2}{[HCl]^4[O_2]}$   
(36) (a)  $Keq = \frac{[HF]^4[O_2]}{[F_2]^2}$ , Keq = 16.2, products (b)  $Keq = \frac{[C_2H_2][H_2]^3}{[CH_4]^2}$  Keq= 0.216, reactants